# Identifying a need for a local infective endocarditis clinic

Alice Jackson<sup>1</sup>, Emma Rose Turner<sup>1</sup>, Caroline Coats<sup>1</sup>, Piotr Sonecki<sup>1</sup>

<sup>1</sup>Queen Elizabeth University Hospital, Glasgow; correspondence to alice.jackson3@nhs.scot

Infective endocarditis (IE) is associated with high levels of morbidity and mortality and

Close surveillance of patients in the first year after an episode of IE, beginning with baseline assessment and repeat echocardiography on completion of antimicrobial therapy, is recommended (2).

In our hospital, which has around 1100 acute adult beds, a weekly IE multidisciplinary team meeting (MDT) facilitates shared care of patients between the cardiology and infectious diseases teams.

Currently, there are no systems in place to ensure timely cardiology clinical and imaging follow-up of patients after an admission with

## **OBJECTIVES**

- To audit current practices in treatment and cardiology follow-up of patients admitted with IE
- To produce an end-of-treatment checklist to incorporate into the IE
- To develop a dedicated IE clinic




### **PRELIMINARY AUDIT**

In Scotland, data hospital admissions are captured through the Scottish Morbidity Scheme (SMR). Consecutive patients with an ICD-10 discharge code for IE (I33, I38, B376) in the first or second diagnostic position from 1/1/18-31/12/18 and 1/1/22-31/12/22 were included. The IE diagnosis was validated. Recurrent hospitalisation and mortality data were linked. Additional clinical data were obtained from electronic records.

In total, 43 patients had a validated discharge diagnosis of IE in the two years reviewed. Mean age was 64 years (IQR 54-77), 58% were men, median length of stay was 40 days (IQR 23-72) and 23% had surgical intervention.

- One-vear all-cause mortality was 37%.
- Overall, 14% of patients were readmitted within 6 weeks (without interim review) and 37% did not have consultant-led cardiology follow-up arranged.
- Median time to outpatient review was 108 days (IQR 53-141) for those who had follow-up arranged and were not readmitted.



- Inpatient death or expected outpatient death
- Readmitted within 6 weeks
- ■None
- Follow-up arranged

### PROGRESS AND IMPLEMENTATION

Audit findings demonstrate that mortality and readmission rates are high after an episode of IE, and patients do not have access to timely cardiology follow-up. It is anticipated that a dedicated IE clinic could lead to improvements in readmission rates and better outcomes for patients.

A proforma containing fields relevant to diagnosis and treatment, as well as specific fields to prompt scheduling of repeat echocardiography and clinic review, has been devised and will be incorporated into routine use at the MDT.

> Plans to pilot a once-monthly, cardiologist-led, one-stop IE clinic, which will combine clinical review with echocardiographic assessment, are ongoing. The clinic will also be used to deliver echocardiography training. Is it anticipated that the IE MDT checklist will trigger referral of those patients who would benefit from early review.

#### **Sta**keholders

**Patients** Consultants **Physiologists** Clinical service managers Cardiology trainees

### **CONCLUSIONS**

There is a need for improved access to timely follow-up for patients after an episode of IE.

To address this, a pilot project is underway to develop a dedicated IE clinic.

Challenges include engaging stakeholders, barriers to introducing a new clinical service while still a trainee, and ensuring access to echocardiographic equipment.

### **REFERENCES**

- (1) Shah ASV, McAllister DA, Gallacher P, et al. Incidence, Microbiology, and Outcomes in Patients Hospitalized With Infective Endocarditis. Circulation. 2020 Jun 23;141(25):2067-2077. doi: 10.1161/CIRCULATIONAHA.119.044913. Epub 2020 May 15. Erratum in: Circulation. 2020 Jun 23;141(25):e960. PMID: 32410460; PMCID: PMC7306256.
- (2) Delgado V, Aimone Marsan N, at al; ESC Scientific Document Group. 2023 ESC Guidelines for the management of endocarditis. Eur Heart J. 2023 Oct 14;44(39):3948-4042. doi: 10.1093/eurhearti/ehad193. Erratum in: Eur Heart J. 2023 Sep 20;: Erratum in: Eur Heart J. 2024 Jan 1;45(1):56.